

On Regular Closure Operators and Cowellpowered Subcategories

Vishvajit V S Gautam

The Institute of Mathematical Sciences, Chennai -600113 India.

vishvajit@imsc.ernet.in

Abstract

Many properties of a category $\mathcal{X}$, as for instance the existence of an adjoint or of a factorization system, are a consequence of the cowellpoweredness of $\mathcal{X}$. In the absence of cowellpoweredness, for general results, fairly strong assumption on the category are needed. This paper provides a number of novel and useful observations to tackle the cowellpoweredness problem of subcategories by means of regular closure operators. Our exposition focusses on the question when two subcategories $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators (up to isomorphism), then information about (non)-cowellpoweredness of $\mathcal{A}$ may be gained from the corresponding property of $\mathcal{B}$, and vice versa.

Keywords : $\mathcal{A}$ -regular morphism, (strongly) epireflective subcategory, cowellpowered category, regular closure operator, (weakly) reflective subcategory, wellpowered category.

AMS subject classification 2000 : 18A20, 18B30, 18A32.

Introduction

Subcategories are always assumed to be full and isomorphism closed. A morphism $f : X \longrightarrow Y$ in a category $\mathcal{X}$ is an *epimorphism* if for each pair of morphisms $g, h : Y \longrightarrow Z$ in a category $\mathcal{X}$ such that $g \cdot f = h \cdot f$ implies $g = h$. Regular closure operators were originally introduced by Salbany [20]. These operators provide a key instrument for attacking the cowellpoweredness problem in a category $\mathcal{X}$. A category $\mathcal{X}$ is said to be *cowellpowered* if each object X in $\mathcal{X}$ admits only a (small) set of non-equivalent $\mathcal{X}$ -epimorphisms with domain in X . If these non-equivalent

$\mathcal{X}$ -epimorphisms form a proper class we say that $\mathcal{X}$ is a non-cowellpowered category. These cases are often settled best by considering the problem in a larger category $\mathcal{B}$ for which the embedding $\mathcal{A} \longrightarrow \mathcal{B}$ preserves epimorphisms. Then clearly $\mathcal{A}$ is cowellpowered whenever $\mathcal{B}$ is.

Using the fully rigid class of topological spaces, Herrlich [10] gave the first example of non-cowellpowered subcategory of **Top**. Schröder [17] produces an example of non-cowellpowered subcategory of **Top**: the full subcategory **Ury** of **Top** having as objects all Urysohn spaces (a topological space X is called *Urysohn* if any pair of distinct points of X can be separated by disjoint closed neighborhoods). Dikranjan, Giuli, Tholen in [4] generalize the Schröder construction [17] and produce more examples of non-cowellpowered subcategories of **HAUS**. In particular, they prove that the category of $S(n)$ -spaces is not cowellpowered. Answering a question whether the category of $S(\omega)$ -spaces is cowellpowered in [4], Dikranjan and Watson [6] proved that for each ordinal $\alpha > 1$ the category of $S(\alpha)$ -spaces is not cowellpowered. It was shown by Giuli and Hušek [8] that the category of topological spaces, in which every compact subspace is Hausdorff, is not cowellpowered. Tozzi [19] proved that the category of topological spaces in which every convergent sequence has a unique limit point is cowellpowered. Some other examples on cowellpoweredness and non-cowellpoweredness can be seen in [2]. A more complete discussion on (non)-cowellpoweredness is given in [5]. See also [7].

A well known technique on the cowellpoweredness problem is to find an effectively defined closure operator c of $\mathcal{X}$ such that the epimorphism of a subcategory $\mathcal{A}$ (whose cowellpoweredness is to be decided) are characterized as c -dense morphism in $\mathcal{A}$. This technique is often difficult because it involves finding an effectively defined closure operator c and then characterizing epimorphisms in terms of c -dense morphism. Schröder remarked in [17] that "Urysohn spaces do not differ very much from Hausdorff spaces, topologically. These small differences imply large consequences. **Ury** is non-cowellpowered and the category **HAUS** of Hausdorff spaces is cowellpowered."

This fact motivated us to study the notion of (non)-cowellpoweredness between two closely related subcategories by means of regular closure operators. In the present paper we discuss various useful constructions for intermediate subcategories, including the Pumplün - Röhrle closure and Hoffmann closure of a subcategory $\mathcal{A}$. Our main focus will be on the question as to when such extensions $\mathcal{B}$ of $\mathcal{A}$ induces the same closure operator as $\mathcal{A}$, so that information on the (non)-cowellpoweredness of $\mathcal{A}$ may be

gained from the corresponding property of $\mathcal{B}$, and vice versa.

Section 1 gives necessary definitions on closure operators. In section 2 we provide the conditions as to when two weakly reflective subcategories induce the same regular closure operators. In section 3 we show that reflective subcategories and their intermediate subcategories induce the same regular closure operators. Section 4 is consisting the main results and presents a necessary and sufficient conditions for two subcategories to induce the same regular closure operators. These results are also obtained for Pumpün-Röhr closure and Hoffmann closure. All the main results on (non)-cowellpoweredness of subcategories are proved in section 5 and examples are given in section 6.

1 Preliminaries

Given two subcategories we give conditions that they induce the same regular closure operators (up to isomorphism). We prove these results for weakly reflective subcategories. These results may be useful in other situation where problem related with cowellpoweredness is not central. Note that, in topological categories regular closure operators are in one to one correspondence with strongly epireflective subcategories (cf. [4]). In section 4 we will give a betterment of this result.

Throughout this paper we consider a category $\mathcal{X}$ and a fixed class $\mathcal{M}$ of morphisms in $\mathcal{X}$ which contains all isomorphisms of $\mathcal{X}$.

It is assumed that $\mathcal{X}$ is $\mathcal{M}$ -complete (cf. [3]), i.e.,

- $\mathcal{M}$ is closed under composition;
- Pullbacks of $\mathcal{M}$ -morphisms exist and belong to $\mathcal{M}$, and multiple pullbacks of (possibly large) families of $\mathcal{M}$ -morphisms with common codomain exist and belong to $\mathcal{M}$.

Some of the consequences of the above assumption are given below.

1. every morphism in $\mathcal{M}$ is a monomorphism in $\mathcal{X}$;
2. if $n \cdot m \in \mathcal{M}$ and $n \in \mathcal{M}$, then $m \in \mathcal{M}$;
3. for each object X in $\mathcal{X}$ the comma category $\mathcal{M}/X$ of $\mathcal{M}$ -morphisms is a (possibly large) complete preordered set. We shall use the usual lattice- theoretical notions in $\mathcal{M}$.

4. there is a (uniquely determined) class $\mathcal{E}$ of $\mathcal{X}$ -morphisms such that $(\mathcal{E}, \mathcal{M})$ is a factorization system of $\mathcal{X}$.

A *closure operator* on $\mathcal{X}$ with respect to the class $\mathcal{M}$ is a family $c = (c_X)_{X \in \mathcal{X}}$ of maps $c_X : \mathcal{M}/X \rightarrow \mathcal{M}/X$ such that

1. $m \leq c(m)$; 2. $m \leq m' \Rightarrow c(m) \leq c(m')$; 3. for every $f : X \rightarrow Y$ and $m \in \mathcal{M}/X$, $f(c_X(m)) \leq c_Y(f(m))$.

Closure operator c determined in this way is unique (up to isomorphism). For each $m \in \mathcal{M}$ we denote by $c(m)$ the *c-closure* of m .

An $\mathcal{M}$ -morphism $m \in \mathcal{M}/X$ is called *c-closed* if $m \cong c_X(m)$. A closure operator c is said to be idempotent if $c(c(m)) \cong c(m)$.

For a subcategory $\mathcal{A}$ of $\mathcal{X}$, a morphism f is an *$\mathcal{A}$ -regular* monomorphism if it is the equalizer of two morphisms $h, k : Y \rightarrow A$ with $A \in \mathcal{A}$.

Let $\mathcal{M}$ contain the class of regular monomorphisms of $\mathcal{X}$. For $m : M \rightarrow X$ in $\mathcal{M}$ define

$$c_{\mathcal{A}}(m) = \wedge \{r \in \mathcal{M} \mid r \geq m \text{ and } r \text{ is } \mathcal{A}\text{-regular}\}$$

These closure operators are called *regular closure operators* and $c_{\mathcal{A}}(m)$ is called the *$\mathcal{A}$ -closure* of m . In case $\mathcal{A} = \mathcal{X}$ we denote $c_{\mathcal{A}}(m)$ by $c(m)$.

Following results are trivial.

1.1. If r is the equalizer of two morphisms f and g , then for every monomorphism α for which $\alpha \cdot f$ and $\alpha \cdot g$ are defined $r = eq(\alpha \cdot f, \alpha \cdot g)$.

1.2. If r is the equalizer of two composite morphisms $\alpha \cdot f$ and $\alpha \cdot g$ with $f \cdot r = g \cdot r$, then $r = eq(f, g)$.

2

We recall the following definitions.

Definition 2.1 A full and replete subcategory $\mathcal{A}$ of a category $\mathcal{X}$ is called weakly reflective ([14]) in $\mathcal{X}$ provided that for each $\mathcal{X}$ -object X there exists a weak $\mathcal{A}$ -reflection, i.e., an $\mathcal{A}$ -object A and a morphism $r : X \rightarrow A$ such that for every morphism $f : X \rightarrow B$ from X into some $\mathcal{A}$ -object B factors through r , i.e., there exists some (not necessarily unique) morphism $g : A \rightarrow B$ with $f = g \cdot r$.

If the reflection morphism r is a monomorphism then we say that $\mathcal{A}$ is weakly monoreflective in $\mathcal{X}$.

Definition 2.2 ([11]) $\mathcal{A}$ is called almost reflective in $\mathcal{X}$ provided that $\mathcal{A}$ is weakly reflective in $\mathcal{A}$ and closed under the formation of retracts in $\mathcal{X}$.

Examples on almost reflective subcategories which are not reflective can be seen in [11]. Clearly, reflectivity $\Rightarrow$ almost reflectivity $\Rightarrow$ weak reflectivity.

Proposition 2.3 Let $\mathcal{A}$ be weakly reflective subcategory of $\mathcal{X}$. If $\mathcal{X}$ has equalizers, cokernel pairs and $\mathcal{M}$ contains all regular monomorphisms of $\mathcal{X}$, then there is a canonical closure operator $c_{\mathcal{A}}$ with respect to $\mathcal{M}$ such that for each $m \in \mathcal{M}$

$$c_{\mathcal{A}}(m) \cong eq(r_Y \cdot i, r_Y \cdot j)$$

with $(i, j : X \longrightarrow Y)$ the cokernel pair of m and r_Y the weak $\mathcal{A}$ -reflection of Y .

Proof. Proof is straightforward and omitted. $\square$

As a consequence of this, $c_{\mathcal{A}}$ is an idempotent closure operators on $\mathcal{X}$.

Assumption: *Throughout the remainder of this paper we assume $\mathcal{X}$ has equalizers, cokernel pairs and $\mathcal{M}$ contains all regular monomorphisms.*


A reflective subcategory of **Top** which contains a space with at least two points has a reflector preserving monomorphism if and only if it is bireflective (i.e., reflection are bijections). But there are categories where notion of monoreflectivity is relevant.

Proposition 2.4 If $\mathcal{A}$ is weakly monoreflective in $\mathcal{X}$, then for each $m \in \mathcal{M}$

$$c_{\mathcal{A}}(m) \cong eq(r_Y \cdot i, r_Y \cdot j) \cong c(m)$$

with $(i, j : X \longrightarrow Y)$ the cokernel pair of m and r_Y the weak $\mathcal{A}$ -reflection of Y .

Proof. Consider the following diagram



with $f = eq(r_Y \cdot i, r_Y \cdot j)$, $m = coker(i, j)$ and $t = eq(p, q)$ a regular monomorphism in $\mathcal{X}$ with $t \geq m$.

Proposition 2.3 gives $f = eq(r_Y \cdot i, r_Y \cdot j) \cong c_{\mathcal{A}}(m)$. Weakly monoreflectivity of $\mathcal{A}$ and 1.1 give $t = eq(r_Z \cdot p, r_Z \cdot q)$, which implies t is also an $\mathcal{A}$ -regular monomorphism. From $f \geq m$ and $t \geq m$ we compute $r_Z \cdot p \cdot m = r_Z \cdot q \cdot m$. Next we use the weakly reflectivity of $\mathcal{A}$ and $m = coker(i, j)$, which give $r_Z \cdot p \cdot f = r_Z \cdot q \cdot f$. Since r_Z is a monomorphism, $r_Z \cdot p \cdot f = r_Z \cdot q \cdot f$ implies $p \cdot f = q \cdot f$. Now $t = eq(p, q)$ and $p \cdot f = q \cdot f$ so there exists a unique h such that $h \cdot t = f$, which implies $f \leq t$. Using the fact that $m \leq f \leq t$ for all regular monomorphism t in $\mathcal{X}$ and the definition of regular closure operators, we deduce that $c_{\mathcal{A}}(m) \cong f \cong c(m)$. $\square$

As a corollary to this we strengthen the result 3.5 of [4].

Corollary 2.5 Any monoreflective $\mathcal{A}$ in $\mathcal{X}$ induces the same regular closure operator as $\mathcal{X}$, which is the largest regular closure operator in $\mathcal{X}$.

Following two observations are the key results for later sections.

Proposition 2.6 Let $\mathcal{A}$ and $\mathcal{B}$ be two weakly reflective subcategories of $\mathcal{X}$. If for each object in $\mathcal{B}$ the weak $\mathcal{A}$ -reflection morphism is a monomorphism, then $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators (up to isomorphism).

Proof. Consider the above diagram (Proposition 2.4) with $f = eq(r_Y \cdot i, r_Y \cdot j) \cong c_{\mathcal{A}}(m)$, $m = coker(i, j)$ and $t = eq(p, q)$ a $\mathcal{B}$ regular monomorphism in $\mathcal{X}$ with $t \geq m$. Since $r_Y \cdot i \cdot m = r_Y \cdot j \cdot m$, there exists a unique k such that $f \cdot k = m$ implies $m \leq f$. $m \leq t$ implies there exists a unique $l : M \rightarrow T$. For $Z \in \mathcal{B}$, r_Z is a monomorphism. Weakly reflectivity of $\mathcal{A}$ gives $t = eq(r_Z \cdot p, r_Z \cdot q)$. Apply l both sides of $r_Z \cdot p \cdot t = r_Z \cdot q \cdot t$ we get $r_Z \cdot p \cdot m = r_Z \cdot q \cdot m$. Since $m = coker(i, j)$, there exists a unique u such that $r_Z \cdot p = u \cdot i$ and $r_Z \cdot q = u \cdot j$. Again apply the

weakly reflectivity of $\mathcal{A}$, we have $v \cdot r_Z = u$. An easy computation gives $r_Z \cdot p \cdot f = r_Z \cdot q \cdot f$, and $t = eq(r_Z \cdot p, r_Z \cdot q)$ implies there exists a unique h such that $t \cdot h = f$ which implies $f \leq t$. Next, using the definition of $c_{\mathcal{B}}(m)$ and the fact that $m \leq c_{\mathcal{A}}(m) \leq t$ for all $\mathcal{B}$ -regular monomorphism t , we prove $c_{\mathcal{A}}(m) \cong c_{\mathcal{B}}(m)$. $\square$

Proposition 2.7 Let $\mathcal{A}$ and $\mathcal{B}$ be weakly reflective subcategories of $\mathcal{X}$. If the composition of weak $\mathcal{B}$ -reflection morphism followed by weak $\mathcal{A}$ -reflection morphism is again an weak $\mathcal{A}$ -reflection morphism, then $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators (up to isomorphism).

Proof. For an object X in $\mathcal{X}$, we denote by r_X the weak $\mathcal{A}$ -reflection of X and by s_X the weak $\mathcal{B}$ -reflection of X .

Consider the diagram

$$\begin{array}{ccccccc}
 M & & & & & & \\
 \downarrow k & \searrow m & & & & & \\
 F & \xrightarrow{f} & X & \xrightleftharpoons[i]{j} & Y & \xrightarrow{s_Y} & sY \xrightarrow{r_{sY}} r(sY) \\
 \downarrow h & \nearrow t & \searrow p & \searrow q & \downarrow w & & \downarrow u \\
 T & & & & Z & \xlongequal{\quad} & Z
 \end{array}$$

with $f = eq(s_Y \cdot i, s_Y \cdot j)$ and $m = coker(i, j)$.

We have $c_{\mathcal{B}}(m) \cong eq(s_Y \cdot i, s_Y \cdot j)$ (Proposition 2.3). Let t be any $\mathcal{A}$ -regular monomorphism with $m \leq t$. Let $t = eq(p, q : X \rightarrow Z)$ with $Z \in \mathcal{A}$. Since $m \leq t$, there exists a unique $l : M \rightarrow T$ such that $t \cdot l = m$. Since $p \cdot m = p \cdot t \cdot l = q \cdot t \cdot l = q \cdot m$, there exists a unique $w : Y \rightarrow Z$ such that $w \cdot i = p$ and $w \cdot j = q$. Now r_{sY} is the weak reflection of Y in $\mathcal{A}$ and $Z \in \mathcal{A}$. Therefore there is a morphism $u : r(sY) \rightarrow Z$ such that $u \cdot r_{sY} \cdot s_Y = w$. By an easy computation we get $p \cdot f = q \cdot f$. As $t = eq(p, q)$, there exists a unique $h : F \rightarrow T$ such that $t \cdot h = f$ which implies $f \leq t$. Since $m \leq f \leq t$ for all $\mathcal{A}$ -regular morphism t , therefore we have $c_{\mathcal{B}}(m) \cong eq(s_Y \cdot i, s_Y \cdot j) \cong c_{\mathcal{A}}(m)$. $\square$

3

In this section we formulate various useful results for intermediate subcategories as to when intermediate subcategory $\mathcal{B}$ of $(\mathcal{A}, \mathcal{X})$ and $\mathcal{A}$ induce

the same regular closure operators (up to isomorphism). Recall that a category $\mathcal{X}$ is said to be wellpowered if the subobjects of every object of $\mathcal{X}$ constitute a (small) set.

Throughout the remainder of this paper by “two subcategories induce the same regular closure operators”, we mean that the induced closure operators are same up to isomorphism (they may be equal).

We begin by defining an intermediate category.

Definition 3.1 [1] Let $\mathbf{r}$ be a reflector from $\mathcal{X}$ to $\mathcal{A}$. If $\mathbf{r} = \mathbf{s} \cdot \mathbf{t}$ where $\mathbf{t} : \mathcal{X} \longrightarrow \mathcal{B}$ is an $\mathcal{X}$ -epireflector and $\mathbf{s} : \mathcal{B} \longrightarrow \mathcal{A}$ is a $\mathcal{B}$ -epireflector, then $\mathcal{B}$ is said to be an *intermediate category* of the pair $(\mathcal{A}, \mathcal{X})$.


Theorem 3.2 Let $\mathcal{X}$ be cowellpowered with products and suppose that every morphism can be factored as an epi followed by mono. Let $\mathcal{A}$ be a reflective subcategory of $\mathcal{X}$ and let $\mathcal{B}$ be a subcategory of $\mathcal{X}$ whose objects are the $\mathcal{X}$ -subobjects of $\mathcal{A}$ -objects. Then $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators.

Proof. We observe that $\mathcal{A}$ is a $\mathcal{B}$ -epireflective subcategory of $\mathcal{B}$ and $\mathcal{B}$ is a $\mathcal{X}$ -epireflective subcategory of $\mathcal{X}$, and that $\mathcal{B}$ is an intermediate category of the pair $(\mathcal{A}, \mathcal{X})$ (cf. [1, Theorem 2]). Now result is immediate from Proposition 2.7. $\square$

Next theorem shows that there exists a smallest intermediate subcategory $\mathcal{B}$ of the pair $(\mathcal{A}, \mathcal{X})$ inducing the same regular closure operators as induced by $\mathcal{A}$.

Theorem 3.3 Let $\mathcal{A}$ be a reflective subcategory of a wellpowered category $\mathcal{X}$ with intersections. The subcategory $\mathcal{B}$ whose objects are the $\mathcal{X}$ -extremal subobjects of $\mathcal{A}$ -objects is the smallest intermediate subcategory of $(\mathcal{A}, \mathcal{X})$ and induces the same regular closure operators as induced by $\mathcal{A}$.

Proof. Consider the diagram



with $f = eq(t_Y \cdot i, t_Y \cdot j)$ and $m = coker(i, j : X \rightarrow Y)$.

Let $n = eq(p, q : X \rightarrow Z)$ be any $\mathcal{A}$ -regular morphism with $n \geq m$. Let $\mathbf{r} : \mathcal{X} \rightarrow \mathcal{A}$ be a $\mathcal{X}$ -reflective functor, $\mathbf{t} : \mathcal{X} \rightarrow \mathcal{B}$ be an $\mathcal{X}$ -epireflective functor and $\mathbf{s} : \mathcal{B} \rightarrow \mathcal{A}$ be a $\mathcal{B}$ -epireflective functor. Using [1, Theorem 5] and Proposition 2.7 we prove that $\mathcal{B}$ is the smallest subcategory of $(\mathcal{A}, \mathcal{X})$ and that $m \leq f \leq n$ for all $\mathcal{A}$ -regular morphisms n . This gives that $c_{\mathcal{B}}(m) \cong f \cong c_{\mathcal{A}}(m)$. $\square$

In the case of largest intermediate subcategory of $(\mathcal{A}, \mathcal{X})$, we have

Theorem 3.4 Let $\mathcal{X}$ be wellpowered, cocomplete category with intersections and products of every indexed set of objects. Let $\mathcal{A}$ be reflective in $\mathcal{X}$ and let $\mathcal{B}$ be the subcategory whose objects are the $\mathcal{X}$ -extremal subobjects of the $\mathcal{A}$ -objects. Define $\mathcal{C}$, a subcategory of $\mathcal{X}$ as follows

$C \in \text{obj}(\mathcal{C}) \iff$ for each $B \in \text{obj}(\mathcal{B})$ and a pair of morphisms $f, g : \mathbf{r}B \rightarrow C$,

$$f \cdot r_B = g \cdot r_B \Rightarrow f = g.$$

(Here, $\mathbf{r}$ is the reflector from $\mathcal{B}$ to $\mathcal{A}$ and $r_B : B \rightarrow \mathbf{r}B$ is the corresponding reflection morphism.) then $\mathcal{C}$ is the largest intermediate subcategory of $(\mathcal{A}, \mathcal{X})$ and induces the same regular closure operators as induced by $\mathcal{A}$.

Proof. (cf. [1, Theorem 6]) We have $\mathcal{C}$ is closed under the formation of products in $\mathcal{X}$ and subobjects in $\mathcal{X}$; $\mathcal{C}$ is thus an epireflective subcategory of $\mathcal{X}$. Let $\mathbf{t} : \mathcal{C} \rightarrow \mathcal{A}$ be the restriction of reflector $\mathbf{s} : \mathcal{X} \rightarrow \mathcal{A}$. It is clear that $\mathbf{t}$ is a reflector. Since $\mathcal{X}$ has (epi, extremal mono)-factorization for morphisms and the way $\mathcal{C}$ was defined, we can prove that for each object C in $\mathcal{C}$, the morphism $\mathbf{t}_C$ is $\mathcal{C}$ -epimorphism and $\mathcal{C}$ is an intermediate subcategory of $(\mathcal{A}, \mathcal{X})$ which is also a largest intermediate subcategory of

$(\mathcal{A}, \mathcal{X})$. Next, use the similar computations as we have done in Theorem 3.3, we get $c_{\mathcal{B}}(m) \cong c_{\mathcal{A}}(m)$. $\square$

4

In this section we give some necessary and sufficient conditions for two subcategories to induce the same regular closure operators.

Throughout this section we assume $\mathcal{X}$ has finite products. ($\mathcal{X}$ also satisfies the assumptions of section 1 and 2).

For a subcategory $\mathcal{A}$ of $\mathcal{X}$, let

$$S(\mathcal{A}) = \{X \in \mathcal{X} \mid \text{there is a monomorphism } X \longrightarrow A \text{ with } A \in \mathcal{A}\}$$

If $\mathcal{A}$ is reflective with reflector $\mathbf{r}$ and reflection morphism r then

$$S(\mathcal{A}) = \{X \in \mathcal{X} \mid r_X \text{ is a monomorphism}\}.$$

If, moreover, $\mathcal{X}$ has (strong epi, mono)-factorization, then $S(\mathcal{A})$ is the *strongly epireflective hull* of $\mathcal{A}$ in $\mathcal{X}$ and $\mathcal{A}$ is bireflective in $S(\mathcal{A})$. For U and X in $\mathcal{X}$,

we denote by

$$\alpha_{U,X} : \mathbf{r}(U \times X) \longrightarrow \mathbf{r}(U) \times \mathbf{r}(X)$$

the *canonical morphism* with $u \cdot \alpha_{U,X} = \mathbf{r}(p)$, $v \cdot \alpha_{U,X} = \mathbf{r}(q)$

where p, q are projections of $U \times X$ and u, v projections of $\mathbf{r}(U) \times \mathbf{r}(X)$.

Let $\triangle_X : X \longrightarrow X \times X$ be the diagonal of X in $\mathcal{X}$.

Theorem 4.1 Let $\mathcal{A}$ and $\mathcal{B}$ be two reflective subcategories of $\mathcal{X}$ and let for each object X in $\mathcal{B}$ the canonical morphism $\alpha_{U,X}$ are defined w.r.t. the reflector $\mathbf{r}$ of the category $\mathcal{A}$ be a monomorphism in $\mathcal{A}$ for all U in $\mathcal{X}$. Then following assertions are equivalent

- (a) $\triangle_X$ is $\mathcal{A}$ -regular for each $X \in \mathcal{B}$;
- (b) $\mathcal{B} \subseteq S(\mathcal{A})$;
- (c) $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators.

Proof. (a) $\Rightarrow$ (b) is immediate from [9, Theorem 1.1]. (b) $\Rightarrow$ (c) follows from Proposition 2.6. (c) $\Rightarrow$ (a) since $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators and $\mathcal{X}$ has cokernel pairs, we have $\triangle_X \cong c_{\mathcal{A}}(\triangle_X) \cong c_{\mathcal{B}}(\triangle_X)$, which completes the result. $\square$

If U is a terminal object, and a generator of $\mathcal{X}$, then the projection $q : U \times X \longrightarrow X$ is an isomorphism, hence also $v \cdot \alpha_{U,X} = r(q)$ is one, so $\alpha_{U,X}$ is monic in $\mathcal{A}$.

Corollary 4.2 Let the terminal object of $\mathcal{X}$ be a generator of $\mathcal{X}$. If $\mathcal{A}$ and $\mathcal{B}$ are reflective subcategories of $\mathcal{X}$, then $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators if and only if $\mathcal{B} \subseteq S(\mathcal{A})$.

Theorem 4.3 of [4] gives that if the terminal object of $\mathcal{X}$ is a generator of $\mathcal{X}$, $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators if and only if $S(\mathcal{A}) = S(\mathcal{B})$. So the Corollary 4.2 is a betterment of this result.

4.3

The Pumplün - Röhl closure $E(\mathcal{A})$ of $\mathcal{A}$.

Let $\mathcal{A}$ be a subcategory of $\mathcal{X}$. A morphism $f : C \rightarrow D$ in $\mathcal{X}$ is called $\mathcal{A}$ -cancellable if $h \cdot f = k \cdot f$ with $h, k : D \rightarrow A$, $A \in \mathcal{A}$ implies $h = k$. Suppose the class $Can_{\mathcal{X}}(\mathcal{A})$ of all $\mathcal{A}$ -cancellable morphisms in $\mathcal{X}$ contains all epimorphisms of $\mathcal{X}$, is closed under composition and colimits, and is stable under (multiple) pushouts. Let $\mathcal{E}$ be a subclass of morphisms of $\mathcal{X}$. An object X in $\mathcal{X}$ is called $\mathcal{E}$ -Hausdorff (cf. [9, 15, 16]) if every $p \in \mathcal{E}$ is $\{X\}$ -cancellable. Let $Haus_{\mathcal{X}}(\mathcal{E})$ be the class of all $\mathcal{E}$ -Hausdorff objects in $\mathcal{X}$ and be closed under mono-sources in $\mathcal{X}$.

We define the Pumplün - Röhl closure $E(\mathcal{A})$ of $\mathcal{A}$ as follows

$$E(\mathcal{A}) = Haus_{\mathcal{X}}(Can_{\mathcal{X}}(\mathcal{A}))$$

Note that $S(\mathcal{A}) \subseteq E(\mathcal{A})$. (cf. [15, 16])

Hoffmann closure $D(\mathcal{A})$ of $\mathcal{A}$

A morphism $f : X \rightarrow Y$ in $\mathcal{X}$ is an $\mathcal{A}$ -epimorphism if for all $u, v : Y \rightarrow A$ with $A \in \mathcal{A}$, one has implication $(u \cdot f = v \cdot f) \Rightarrow (u = v)$. This definition is same as the definition of $\mathcal{A}$ -cancellable morphism, but we reserve their names as many other authors do.

Let $Epi_{\mathcal{X}}(\mathcal{A})$ be the class of all $\mathcal{A}$ -epimorphisms of $\mathcal{X}$. We denote by

$$D(\mathcal{A}) = Haus_{\mathcal{X}}(Epi_{\mathcal{X}}(\mathcal{A}))$$

the Hoffmann closure of $\mathcal{A}$. One has $\mathcal{A} \subseteq S(\mathcal{A}) \subseteq E(\mathcal{A}) \subseteq D(\mathcal{A}) \subseteq \mathcal{X}$ ([9]), and all inclusions may be proper. If $\mathcal{C}$ is one of these three intermediate categories, then

- (1) $\mathcal{A} \rightarrow \mathcal{C}$ preserves epimorphisms,
- (2) $\mathcal{A}$ is epireflective in $\mathcal{C}$ if $\mathcal{A}$ is reflective in $\mathcal{C}$,
- (3) $\mathcal{C}$ is strongly reflective in $\mathcal{X}$ if $\mathcal{X}$ has (strong epi, mono)-factorization.

Every morphism in $(m \mid p \perp m \text{ for all } p \in Epi_{\mathcal{X}}(\mathcal{A}))$ is a strong monomorphism and is called $\mathcal{A}$ -straight in [9]. $\mathcal{A}$ -strong monomorphisms are $\mathcal{A}$ -straight.

Theorem 4.4 Let $\mathcal{A}$ and $\mathcal{B}$ be reflective subcategories of $\mathcal{X}$. Consider the following assertions:

- (a) $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators ;
- (b) $\mathcal{B} \subseteq E(\mathcal{A})$ (if every epimorphism in $\mathcal{X}$ is $\mathcal{A}$ -cancellable);
- (c) $\mathcal{B} \subseteq D(\mathcal{A})$.

Then (a) $\Rightarrow$ (b) $\Rightarrow$ (c).

Moreover, if the terminal object of $\mathcal{X}$ is a generator, then

(a) $\Leftrightarrow$ (b) and (a) $\Leftrightarrow$ (c).

Proof. (a) $\Rightarrow$ (b). For every object $X \in \mathcal{X}$, we have Δ_X is $\mathcal{A}$ -strong implies $X \in E(\mathcal{A})$ (cf. [9, Corollary 2.2]). $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators, for each $X \in \mathcal{B}$ we have Δ_X is $\mathcal{A}$ -regular. In $\mathcal{X}$ every $\mathcal{A}$ -regular monomorphism is $\mathcal{A}$ -strong monomorphism. This shows $\mathcal{B} \subseteq E(\mathcal{A})$.

(b) $\Rightarrow$ (c) We have $\mathcal{B} \subseteq E(\mathcal{A}) \subseteq D(\mathcal{A})$.

Moreover part

(b) $\Rightarrow$ (a) We have $\mathcal{B} \subseteq E(\mathcal{A})$. Corollary 4.2 gives $\mathcal{B} \subseteq S(\mathcal{A})$, We also have $S(\mathcal{A}) \subseteq E(\mathcal{A})$. Now the result follows from the fact that $\mathcal{B} \subseteq S(\mathcal{A}) \subseteq E(\mathcal{A})$.

(c) $\Rightarrow$ (a) Corollary 4.2 gives $\mathcal{B} \subseteq S(\mathcal{A})$, the result is immediate from the fact that $S(\mathcal{A}) \subseteq D(\mathcal{A})$ and $\mathcal{B} \subseteq S(\mathcal{A}) \subseteq D(\mathcal{A})$. $\square$

In case of intermediate subcategories, we have

Theorem 4.5 Let $\mathcal{X}$ be cowellpowered and has (epi, mono)- factorizations. Let $\mathcal{A}$ be a reflective and cowellpowered subcategory of $\mathcal{X}$. Let $\mathcal{B}$ be a subcategory of $\mathcal{X}$ such that for each object X in $\mathcal{B}$ the canonical morphism $\alpha_{U,X}$ are defined w.r.t. the reflector $\mathbf{r}$ of the category $\mathcal{A}$ is a monomorphism in $\mathcal{A}$ for all U in $\mathcal{X}$. Then

(a) if objects of $\mathcal{B}$ are the $\mathcal{X}$ -subobjects of $\mathcal{A}$ -objects, then the $\mathcal{B} \subseteq S(\mathcal{A})$.

(b) if objects of $\mathcal{B}$ are the $\mathcal{X}$ -extremal subobjects of $\mathcal{A}$ -objects, then $\mathcal{B}$ is smallest intermediate subcategory of $(\mathcal{A}, \mathcal{X})$ contained in $S(\mathcal{A})$.

Moreover, if the terminal object of $\mathcal{X}$ is a generator, then the results hold without the assumption that $\alpha_{U,X}$ is a monomorphism in $\mathcal{A}$ for all U in $\mathcal{X}$.

In addition to this, if $\mathcal{X}$ has (strong epi, mono)-factorizations, $\mathcal{B} = S(\mathcal{A}) = S(\mathcal{B})$.

Proof. (a) Since $\mathcal{X}$ is cowellpowered and has (epi, mono)-factorization and $\mathcal{B}$ forms an intermediate subcategory of the pair $(\mathcal{A}, \mathcal{X})$ such that $\mathcal{A}$ is

$\mathcal{B}$ -epireflective subcategory of $\mathcal{B}$ and $\mathcal{B}$ is $\mathcal{X}$ -epireflective subcategory of $\mathcal{X}$. Therefore $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators (cf. Theorem 3.2). Now result follows from Theorem 4.1.

(b) Proof is clear from Theorems 3.3 , 4.1. $\square$

Let $\mathcal{B}$ be any reflective subcategory of $\mathcal{X}$, suppose $\mathcal{A}$ be another reflective subcategory of $\mathcal{X}$. One can ask, what will be the suitable conditions on $\mathcal{A}$ such that $\mathcal{B} \subseteq \mathcal{A}$? Following theorem provides an answer to this question by means of regular closure operators.

Theorem 4.6 Let $\mathcal{B}$ be any reflective subcategory of $\mathcal{X}$. If $\mathcal{A}$ is an extremally epireflective subcategory of $\mathcal{X}$ and X is not an object of $\mathcal{A}$, then there exist morphisms $f, g : Z \longrightarrow X$ such that $(*) f \neq g$ and $r_X \cdot f = r_X \cdot g$, and a morphism $\bar{g} : X \longrightarrow Z$ such that $\bar{g} \cdot g = \bar{g} \cdot f$, and $g \cdot \bar{g} \cdot g = g$, where r_X is the $\mathcal{A}$ -reflection morphism. If $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators then $\mathcal{B} \subseteq \mathcal{A}$.

Proof. Suppose $X \in \mathcal{B}$, this implies Δ_X is $\mathcal{B}$ -regular. Since $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators, $\Delta_X \cong c_{\mathcal{A}}(\Delta_X) \cong c_{\mathcal{B}}(\Delta_X)$. This implies Δ_X is $\mathcal{A}$ -regular. Since $X \in \mathcal{A}$ if and only if Δ_X is $\mathcal{A}$ -regular, $X \in \mathcal{A}$ (cf. [13, Proposotion 3.4]). Hence $\mathcal{B} \subseteq \mathcal{A}$.

Remark.(cf. [13]) Some examples of categories $\mathcal{X}$ in which condition $(*)$ is satisfied are: all topological categories; many initially structured categories; such as **HAUS**; and the category of groups.

5

Cowellpoweredness - main results

In this section (except in the Theorem 5.2) category $\mathcal{X}$ is assumed to be complete, wellpowered and cowellpowered.

Theorem 5.1 Let $\mathcal{A}$ be reflective and cowellpowered subcategory of $\mathcal{X}$. Let $\mathcal{B}$ be a reflective subcategory of $\mathcal{X}$, consider the following conditions on $\mathcal{A}$ and $\mathcal{B}$.

- (a) for each object X in $\mathcal{B}$ the canonical morphism $\alpha_{U,X}$ is a monomorphism in $\mathcal{A}$ for all U in $\mathcal{X}$.
- (b) objects of $\mathcal{B}$ are the $\mathcal{X}$ -subobjects of $\mathcal{A}$ -objects.
- (c) objects of $\mathcal{B}$ are the $\mathcal{X}$ -extremal subobjects of $\mathcal{A}$ -objects.

- (d) the terminal object of $\mathcal{X}$ is a generator of $\mathcal{X}$.
- (e) $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators.
- (f) $\mathcal{B}$ is cowellpowered.
- (g) $\mathcal{B}$ is an intermediate cowellpowered subcategory of $(\mathcal{A}, \mathcal{X})$.
- (h) $\mathcal{B} = S(\mathcal{A}) = S(\mathcal{B})$ is cowellpowered.

Then $((a) \text{ and } (e)) \Rightarrow (f) \Leftarrow ((d) \text{ and } (e)); ((a) \text{ and } (b)) \Rightarrow (g) \Leftarrow ((b) \text{ and } (d))$ and $((a) \text{ and } (c)) \Rightarrow (h) \Leftarrow ((c) \text{ and } (d))$.

Proof. $((a) \text{ and } (e)) \Rightarrow (f)$. Theorem 4.1 gives $\mathcal{B} \subseteq S(\mathcal{A})$. Since $\mathcal{A}$ is reflective and cowellpowered, $S(\mathcal{A})$ is cowellpowered (cf. [12]) and hence $\mathcal{B}$ is cowellpowered.

$((d) \text{ and } (e)) \Rightarrow (f)$. $\mathcal{X}$ is complete and wellpowered therefore $\mathcal{X}$ has (strong epi, mono)-factorizations. Since $\mathcal{A}$ and $\mathcal{B}$ are reflective in $\mathbf{X}$, therefore $S(\mathcal{A})$ and $S(\mathcal{B})$ are the strongly epireflective hulls of $\mathcal{A}$ and $\mathcal{B}$ respectively. By the fact that $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators we have $S(\mathcal{A}) = S(\mathcal{B})$. $\mathcal{A}$ is also cowellpowered in $\mathcal{X}$ therefore we have $S(\mathcal{A})$ is cowellpowered and hence $\mathcal{B}$ is cowellpowered.

$((a) \text{ and } (b)) \Rightarrow (g)$. Proof follows from the facts that $\mathcal{B} \subseteq S(\mathcal{A})$ (cf. Theorem 4.5) and $S(\mathcal{A})$ is cowellpowered (cf. [12], [18]).

$((b) \text{ and } (d)) \Rightarrow (g)$. Clearly, $S(\mathcal{A}) = S(\mathcal{B})$. Now the result is immediate from the fact that $S(\mathcal{A})$ is cowellpowered.

$((a) \text{ and } (c)) \Rightarrow (h) \Leftarrow ((c) \text{ and } (d))$. Left for reader. $\square$

We conclude this section with the following result (cf. Theorem 4.6).

Theorem 5.2 Let $\mathcal{A}$ be extremally epireflective subcategory of $\mathcal{X}$ satisfying condition (*) of Theorem 4.6. Let $\mathcal{B}$ be a non-cowellpowered reflective subcategory of $\mathcal{X}$ such that $\mathcal{B} \longrightarrow \mathcal{A}$ preserves epimorphisms. If $\mathcal{A}$ and $\mathcal{B}$ induce the same regular closure operators, then $\mathcal{A}$ is noncowellpowered.

Proof. Proof is a straightforward consequence of Theorem 4.6. $\square$

6

Examples

(1) (cf. [1]) The category **CHS** of compact Hausdorff spaces is a reflective subcategory of **Top** and the category **CRS** of completely regular spaces form the smallest intermediate subcategory for the pair $(\mathbf{CHS}, \mathbf{Top})$ whose objects are the subspaces of the objects of **CHS** and the reflection of an object X is the Stone-Cech compactification of X . So

CHS and **CRS** induce the same regular closure operators on **Top**. Also, **CHS** form the smallest monoreflective subcategory of **CRS**, therefore for each $X \in \mathbf{CRS}$ and $M \subseteq X$, $c_{\mathbf{CHS}}(M) = c_{\mathbf{CRS}}(M)$.

(2) (cf. [1]) Let **Top_o** denote the category of T_o -spaces. Let $\mathcal{A}$ be a subcategory of **Top_o** whose objects are **Top_o**- extremal subobjects of products of the connected doublet (the two point space with three open sets). This category $\mathcal{A}$ is the smallest monoreflective subcategory of **Top_o**. Therefore $\mathcal{A}$ and **Top_o** induce the same regular closure operators.

Another example of this fact is the category of indiscrete spaces which forms the smallest monoreflective subcategory of **Top**.

(3) (cf. [2]) Let $\mathcal{P}$ be a class of topological spaces and let **HAUS**($\mathcal{P}$) denote the category of topological spaces X such that for every $P \in \mathcal{P}$ and for every continuous map $f : P \rightarrow X$, $f(P)$ is a Hausdorff subspace of X . Let $\mathcal{P}$ satisfy the following condition:

If $P \in \mathcal{P}$, $P \neq \emptyset$ then $P \cup Q \in \mathcal{P}$ for some non- empty space Q and $\mathcal{P}$ is closed under continuous images.

Since **HAUS**($\mathcal{P}$) and **HAUS**($\mathcal{P} \cap \mathbf{HAUS}$) induce the same regular closure operators (cf. [2, Theorem 1.4]), we have **HAUS**($\mathcal{P}$) $\subseteq$ **HAUS**($\mathcal{P} \cap \mathbf{HAUS}$) (cf. Corollary 4.2). Now the non- cowellpoweredness of **HAUS**($\mathcal{P} \cap \mathbf{HAUS}$) follows from the non-cowellpoweredness of **HAUS**($\mathcal{P}$) and the cowellpoweredness of **HAUS**($\mathcal{P}$) follows from the cowellpoweredness of **HAUS**($\mathcal{P} \cap \mathbf{HAUS}$) (cf. Theorem 5.1). In particular non- cowellpoweredness of **HAUS**(**comp**) ([8, Theorem 4.3]) implies the non-cowellpoweredness of **HAUS**(**Hcomp**), where **comp** denotes the class of all compact spaces and **Hcomp** denotes the class of all compact Hausdorff spaces.

Acknowledgements: I am thankful to the referee for his valuable and helpful suggestions.

I wish to thank Professor Walter Tholen for his very constructive comments of an earlier version of this paper. I also wish to express my thanks to Professor Parameswaran Sankaran, IMSc Chennai, for his continuous encouragement.

References.

- [1] S. Baron, Reflectors as compositions of epireflectors, Trans. Amer. Math. Soc. 136(1969) 499-508.
- [2] D. Dikranjan and E. Giuli, Ordinal invariants and epimorphisms in some categories of weak Hausdorff spaces, Comment. Math. Univ. Carolin. 27(1986) 395-417.
- [3] D. Dikranjan and E. Giuli, Closure operators I, Topology Appl. 27(1987) 129-143.

- [4] D. Dikranjan, E. Giuli and W. Tholen, Closure operators II, in: Categorical topology and its relation to Analysis, Algebra and Combinatorics, Proceedings International Conference Prague, 1988 (World Scientific, Singapore, 1989) 297-335.
- [5] D. Dikranjan and W. Tholen, Categorical structure of closure operators with applications to Topology, Algebra and Discrete Mathematics (Kluwer, Dordrecht, 1994).
- [6] D. Dikranjan and S. Watson, The category of $S(\alpha)$ -spaces is not cowellpowered, Topology Appl. 61(1995) 137-150.
- [7] V.V.S. Gautam, On some aspects of Banaschewski-Fomin-Shanin extension and closure operators, Ph.D. Thesis, Kurukshetra University, 1998.
- [8] E. Giuli and M. Hušek, A diagonal theorem for epireflective subcategories of **Top** and cowellpoweredness, Ann. Mat. Pura Appl. (4) 145(1986) 337-346.
- [9] E. Giuli, S. Mantovani and W. Tholen, Objects with closed diagonals, J. Pure Appl. Algebra 51(1988) 129-140.
- [10] H. Herrlich, Epireflective subcategories of **Top** need not be cowellpowered, Comment. Math. Univ. Carolin. 16(1975) 713-715.
- [11] H. Herrlich, Almost reflective subcategories of **Top**, Topology Appl. 49(1993) 251-264.
- [12] R.E. Hoffmann, Cowellpowered reflective subcategories, Proc. Amer. Math. Soc. 90(1984) 45-46.
- [13] H. Lord, Factorization, diagonal separation, and disconnectedness, Topology Appl. 47(1992) 83-96.
- [14] P.C. Kainen, Weak adjoint functors, Math. Z. 122(1971) 1-9.
- [15] D. Pumplün, Die Hausdorff-Korrespondenz, unpublished manuscript, 14pp, Hagen, 1976.
- [16] D. Pumplün and H. Röhl, Separated totally convex spaces, Manuscripta Math. 50(1985) 145-183.
- [17] J. Schröder, The category of Urysohn spaces is not cowellpowered, Topology Appl. 16(1983) 237-241.
- [18] W. Tholen, Reflective subcategories, Topology Appl. 27(1987) 201-212.
- [19] A. Tozzi, US-spaces and closure operators, Rend. Circ. Mat. Palermo(2) Suppl. 12(1986) 291-300.
- [20] S. Salbany, Reflective subcategories and closure operators, in : Lecture Notes in Mathematics 540(Springer, Berlin, 1976) 548-565.